

Last time $E(XY) = E(X)E(Y)$ if X, Y indep.

Next, $\text{Var}(X+Y) = \text{Var}(X) + \text{Var}(Y)$ if X, Y indep

Recall, $\text{Var}(X) = E((X - E(X))^2) = E((X - \mu_X)^2)$
 $\mu_X = E(X)$.

$$\begin{aligned} \text{Var}(X+Y) &= E\left(\left((X+Y) - E(X+Y)\right)^2\right) \\ &= E\left(\left((X+Y) - (\mu_X + \mu_Y)\right)^2\right) \\ &= E\left(\left((X - \mu_X) + (Y - \mu_Y)\right)^2\right) \end{aligned}$$

FOIL

$$= E\left(\left(X - \mu_X\right)^2 + 2(X - \mu_X)(Y - \mu_Y) + (Y - \mu_Y)^2\right)$$

$$= E\left(\left(X - \mu_X\right)^2\right) + 2E\left(\left(X - \mu_X\right)(Y - \mu_Y)\right) + E\left(\left(Y - \mu_Y\right)^2\right)$$

"
Var(X)

$$2E(X - \mu_X)E(Y - \mu_Y)$$

"
Var(Y)

indep.
of X, Y

"
0

Note $E(X - \mu_X) = E(X) - \mu_X = \mu_X - \mu_X = 0$

So $\text{Var}(X+Y) = \text{Var}(X) + \text{Var}(Y)$ when X, Y indep.

(2)

$$\text{ex } X \sim \text{Bin}(n, p)$$

$X = I_1 + \dots + I_n$ sum of indep Bernoulli trials,

$$E(X) = E(I_1) + \dots + E(I_n) = np$$

" " "

p p

$$\text{Var}(X) = \text{Var}(I_1) + \dots + \text{Var}(I_n) = np(1-p)$$

" "

p(1-p)

$$\text{SD}(X) = \sqrt{npq}$$

Expectation and SD of Sample sum and average

X_1, X_2, \dots, X_n iid $E(X_i) = \mu$, $\text{SD}(X_i) = \sigma$

let $S_n = X_1 + \dots + X_n$ sample sum,

$$E(S_n) = n\mu \quad (\text{identically distributed})$$

$$\text{SD}(S_n) = \sqrt{n\sigma^2} = \sqrt{n}\sigma \quad \text{Var}(S_n) = n\sigma^2$$

let $A_n = \frac{S_n}{n}$ sample avg.

$$E(A_n) = E\left(\frac{S_n}{n}\right) = \frac{E(S_n)}{n} = \mu$$

$$\text{SD}(A_n) = \text{SD}\left(\frac{S_n}{n}\right) = \frac{1}{n} \text{SD}(S_n) = \frac{1}{n} \sqrt{n}\sigma = \frac{\sigma}{\sqrt{n}}$$


Central Limit Theorem (CLT)

X_1, \dots, X_n iid mean= μ , SD= σ

$S_n = X_1 + \dots + X_n, A_n = \frac{S_n}{n}$



Then for "large" n , the distribution of S_n and A_n are normal,

Picture



draw n w/ replacement X_1, X_2, \dots, X_n

$S_n = X_1 + X_2 + \dots + X_n$



For CLT app:
Google Brown CLT Seeing Theory

ex (3.3.17)

Let X be a RV w/ $P(X=-100) = P(X=0) = 1/4, P(X=100) = 1/2$

Let S be sum of 25 indep RV each with the same dist as X .


Find $P(S=0)$

$$\begin{array}{ccc} & 1 & 1 & 2 \\ & / & / & / \\ \boxed{-100} & 0 & 100 \end{array}$$

draw 25
 X_1, X_2, \dots, X_{25}

$$S = X_1 + \dots + X_{25}$$

Values
 -2500 to 2500
 increment of
 100
 ← bins
 width
 100.



$$E(X) = -100 \cdot \frac{1}{4} + 0 \cdot \frac{1}{4} + 100 \cdot \frac{1}{2}$$

$$= 25$$

$$SD(X) = \sqrt{E(X^2) - E(X)^2}$$

$$E(X^2) = (-100)^2 \cdot \frac{1}{4} + 0^2 \cdot \frac{1}{4} + (100)^2 \cdot \frac{1}{2}$$

$$E(X)^2 = 25^2$$

$$SD(X) = 100 \frac{\sqrt{11}}{4}$$

$$E(S) = 25^2$$

$$SD(S) = 100 \frac{\sqrt{11}}{4} \cdot 5$$

$$\Phi\left(\frac{50 - 25^2}{\frac{500\sqrt{11}}{4}}\right) - \Phi\left(\frac{-50 - 25^2}{\frac{500\sqrt{11}}{4}}\right) = 0.031$$

Finished with section 3.3,

Next sec 3.6. See Reading Guide what to read in this section,

Understand how to find variance of a sum of non independent indicators as in next example,

Stat 134

Friday February 16 2018

1. A building has 10 floors above the basement. If 12 people get into an elevator at the basement, and each chooses a floor at random to get out, independently of the others, at how many floors do you expect the elevator to make a stop to let out one or more of these 12 people?

a $12(1 - (9/10)^{10})$

b $10(1 - (9/10)^{12})$

c none of the above

$$E(X) = 1 - \left(\frac{9}{10}\right)^{12}$$

$X = \#$ elevator stops

$$X = I_1 + \dots + I_{10}$$

$$E(X) = 10 \left(1 - \left(\frac{9}{10}\right)^{12}\right)$$


$I_i =$ indicator for at least 1 person gets off at floor i

Find $Var(X)$.

$Var(X) = Var(I_1) + \dots + Var(I_{10})$ since I_j not indep.

$$E(X^2) = E\left(\left(\sum_{j=1}^{10} I_j\right)^2\right)$$

$$\begin{aligned} & \left(\sum_{j=1}^{10} I_j\right)^2 \\ &= \sum_{j=1}^{10} I_j^2 + \sum_{1 \leq i \neq j \leq 10} 2I_i I_j \end{aligned}$$



$I_1 I_2$ = indicator - stop at 1st and 2nd floor

I_{12} = at least one person gets off at 1st and 2nd floor

$E(I_{12}) = 1 - \text{Prob}(\text{no one gets off 1st floor or no one gets off 2nd floor})$

$$\begin{aligned} & \text{Prob}(\text{no one at 1st floor}) + \text{Prob}(\text{no one at 2nd floor}) - \text{Prob}(\text{no one at 1st and 2nd floor}) \\ & \left(\frac{9}{10}\right)^{12} + \left(\frac{9}{10}\right)^{12} - \left(\frac{8}{10}\right)^{12} \\ & \left(\frac{8}{10}\right)^{12} \end{aligned}$$

$$\begin{aligned} \underline{E(x^2)} &= E\left(\sum_{j=1}^{10} I_j\right) + E\left(\sum_{1 \leq i < j \leq 10} I_i I_j\right) \\ &= n E(I_1) + 10 \cdot 9 E(I_1 I_2) \end{aligned}$$

$$\underline{\text{Var}(x)} = E(x^2) - E(x)^2 = \boxed{.34}$$

Next understand derivation of variance of hypergeometric distribution. I worked out the details for you on the next page.

Variance of hypergeometric (N, G, n) Prob = G/N

$$X = I_1 + \dots + I_n \quad \text{where } I_i = \begin{cases} 1 & \text{if } i^{\text{th}} \text{ element is Good.} \\ 0 & \text{else.} \end{cases}$$

$$\text{Know } E(X) = n \left(\frac{G}{N} \right)$$

$$E(X^2) = E\left(\left(\sum_{j=1}^n I_j\right)^2\right) = E\left(\sum_{j=1}^n I_j^2 + \sum_{i \neq j} I_i I_j\right)$$

$$= nE(I_1) + n(n-1)E(I_1 I_2)$$

$$I_1 I_2 = I_{12} = \begin{cases} 1 & \text{if } 1^{\text{st}} \text{ and } 2^{\text{nd}} \text{ element is Good} \\ 0 & \text{else} \end{cases} \quad \leftarrow \text{Prob } \frac{G}{N} \frac{G-1}{N-1}$$

$$\Rightarrow E(X^2) = n \frac{G}{N} + n(n-1) \frac{G}{N} \frac{G-1}{N-1}$$

$$\Rightarrow \text{Var}(X) = E(X^2) - E(X)^2 = n \frac{G}{N} + n(n-1) \frac{G}{N} \frac{G-1}{N-1} - \left(n \frac{G}{N}\right)^2$$

$$= \frac{nG}{N} \left[1 + \frac{(n-1)(G-1)}{(N-1)} - \frac{nG}{N} \right]$$

$$= \frac{nG}{N} \left[\frac{N(N-1) + (n-1)(G-1)N - nG(N-1)}{N(N-1)} \right]$$

$$= \frac{nG}{N} \left[\frac{N^2 - \cancel{nN} + \cancel{nGN} - \cancel{GN} - \cancel{nN} + \cancel{nGN} - \cancel{nGN} + nG}{N(N-1)} \right]$$

$$= \frac{nG}{N} \left[\frac{N^2 - 6N - nN + nG}{N(N-1)} \right]$$

$$= \frac{nG}{N} \left[\frac{N(N-6) - n(N-G)}{N(N-1)} \right]$$

$$= \frac{nG}{N} \left[\frac{(N-6)(N-n)}{N \cdot N-1} \right]$$

$$= \frac{n}{N} \left(\frac{G}{N} \right) \left(\frac{N-6}{N} \right) \left(\frac{N-n}{N-1} \right)$$

$$\Rightarrow \boxed{\text{SD}(X) = \sqrt{n \left(\frac{G}{N} \right) \left(1 - \frac{G}{N} \right)} \cdot \sqrt{\frac{N-n}{N-1}}}$$