

Warmup 11:00 - 11:10

Let (X, Y) be bivariate normal. Then $(2X+3Y+4, 6X-Y-4)$ is bivariate normal.

a true

b false

c not enough info to decide

Show $a(2X+3Y+4) + b(6X-Y-4)$ is normal for all a, b .

$$= \underbrace{(2a+6b)X + (3a-b)Y}_{\substack{\uparrow \\ \text{normal since} \\ (X, Y) \text{ BUN}}} + \underbrace{4a-4b}_{\substack{\uparrow \\ \text{constant}}} \text{ is normal}$$

$\Rightarrow (2X+3Y+4, 6X-Y-4)$ is BUN

Announcement:

Review materials are on stat134.org website.

Total Variance Decomposition

Sec 6.4

Properties (from lec 34)


- ① $E(Y) = E(E(Y|X))$ iterated expectation
- ② $E(aY+b|X) = aE(Y|X) + b$
- ③ $E(Y+Z|X) = E(Y|X) + E(Z|X)$
- ④ $E(g(X)|X) = g(X)$
- ⑤ $E(g(X)Y|X) = g(X)E(Y|X)$
- ⑥ $Var(Y) = E(Var(Y|X)) + Var(E(Y|X))$ total variance decomposition (sec 6.2.18)

Lab 1	Lab 2	Lab 3	Lab 4	Lab 5	Lab 6	Lab 7
4.13	3.86	4.00	3.88	4.02	4.02	4.00
4.07	3.85	4.02	3.88	3.95	3.86	4.02
4.04	4.08	4.01	3.91	4.02	3.96	4.03
4.07	4.11	4.01	3.95	3.89	3.97	4.04
4.05	4.08	4.04	3.92	3.91	4.00	4.10
4.04	4.01	3.99	3.97	4.01	3.82	3.81
4.02	4.02	4.03	3.92	3.89	3.98	3.91
4.06	4.04	3.97	3.90	3.89	3.99	3.96
4.10	3.97	3.98	3.97	3.99	4.02	4.05
4.04	3.95	3.98	3.90	4.00	3.93	4.06

total variance is the average of the group variances (written $E(Var(Y|X))$) plus the variance of the group averages (written $Var(E(Y|X))$).

This is proved in exercise 6.7.18.

Picture



ex Let $U \sim \text{Unif}(0,1)$ and $X|U \sim \text{Exp}\left(\frac{1}{U}\right)$

Recall if $E\left(\text{Exp}\left(\frac{1}{U}\right)\right) = U \Rightarrow E(X|U) = U$

$\text{Var}\left(\text{Exp}\left(\frac{1}{U}\right)\right) = U^2 \Rightarrow \text{Var}(X|U) = U^2$

Find $E(X)$, $E(UX)$ and $\text{Var}(X)$.

$$E(X) = E(E(X|U)) = E(U) = \left(\frac{1}{2}\right)$$

$$\begin{aligned} E(UX) &= E(E(UX|U)) = E(U \underbrace{E(X|U)}_U) \\ &= E(U^2) = \underbrace{\text{Var}(U)}_{\frac{1}{12}} + \underbrace{E(U)^2}_{\left(\frac{1}{2}\right)^2} = \left(\frac{1}{3}\right) \end{aligned}$$

$$\text{Var}(X) = \underbrace{E(\underbrace{\text{Var}(X|U)}_{U^2})}_{\frac{1}{3}} + \underbrace{\text{Var}(E(X|U))}_0 = \left(\frac{5}{12}\right)$$



Joy Lin

1:11pm


HW 13, 6.3.12, b)

6.rev.21, c)

12. Suppose there are ten atoms, each of which decays by emission of an α -particle after an exponentially distributed lifetime with rate 1, independently of the others. Let T_1 be the time of the first α -particle emission, T_2 the time of the second. Find:

- a) the distribution of T_1 ;
- b) the conditional distribution of T_2 given T_1 ;
- c) the distribution of T_2 .

a) $T_1, \dots, T_{10} \stackrel{iid}{\sim} \text{exp}(1)$



$$P(T_{(1)} \in dx) = \binom{10}{1, 9} 1 \cdot e^{-1 \cdot x} \cdot (e^{-x})^9 dx = \boxed{10e^{-10x} dx}$$



$$P(T_{(1)} \in dx, T_{(2)} \in dy) = \binom{10}{1, 8} 1 e^{-x} \cdot 8 e^{-y} (e^{-y})^8$$

$$= 10 \cdot 8 e^{-x-y}$$

$$\Rightarrow P(T_{(2)} \in dy | T_{(1)} \in dx) = \frac{P(T_{(1)} \in dx, T_{(2)} \in dy)}{P(T_{(1)} \in dx)}$$

$$= \frac{80 e^{-x-y}}{10 e^{-10x}} \frac{dy}{dx} = \boxed{8 e^{-9(y-x)} dy}$$

$$\begin{aligned}
 c) P(T_{(2)} \leq y) &= \int_{x=0}^y 9e^{-9(y-x)} \cdot 10e^{-10x} dx dy \\
 &= 90 \int_{x=0}^y e^{-9y} e^{xy} e^{-10x} dx dy = \boxed{90e^{-9y}(1-e^{-y}) dy}
 \end{aligned}$$

21. I toss a coin which lands heads with probability p . Let W_H be the number of tosses till I get a head, W_{HH} the number of tosses till I get two heads in a row, and W_{HHH} the number of tosses till I get three heads in a row. Find:

- a) $E(W_H)$; b) $E(W_{HH})$ [Hint: condition on whether the first toss was heads or tails]; c) $E(W_{HHH})$ [Hint: condition on W_T].
 d) Generalize to find the expected number of tosses to obtain m heads in a row.

better to condition on W_{HH}

$$\begin{aligned}
 E(W_{HHH}) &= E(E(W_{HHH} | W_{HH})) \\
 &= \sum_{s=1}^{\infty} E(W_{HHH} | W_{HH} = s) \cdot P(W_{HH} = s)
 \end{aligned}$$

let $\mu = E(W_{HHH})$

W_{HH}	$P(W_{HH})$	$E(W_{HHH} W_{HH})$	
2	p^2	$(\mu+3)p + 3p$	AHT, HHT
3	$2p^2$	$(\mu+4)p + 4p$	THT, THT, THH, THH
4	$2^2 p^2$	$(\mu+5)p + 5p$	TTHT, TTHT, TTH, TTH
...	...		

$$\begin{aligned}
\mu &= \sum_{x=2}^{\infty} [(x+1)q + (x+1)p] P(W_{HH} = x) \\
&= \sum_{x=2}^{\infty} [(x+1) + q\mu] P(W_{HH} = x) \\
&= E(W_{HH} + 1) + \mu q \\
&= \frac{1}{p} + \frac{1}{p^2} + 1 + \mu q
\end{aligned}$$

$$\Rightarrow p\mu = 1 + \frac{1}{p} + \frac{1}{p^2}$$

$$\Rightarrow \boxed{\mu = \frac{1}{p} + \frac{1}{p^2} + \frac{1}{p^3}}$$



Ksenia Bogdanova

6:36pm

HW 11, 5.rev.24 (c and d)

24. A coin of diameter d is tossed at random on a grid of squares of side s . Making appropriate assumptions, to be stated clearly, calculate:

a) the probability that the coin lands inside some square (i.e., not touching any line);


b) the probability that the coin lands heads inside some square.

Suppose now that the coin is tossed four times. Let X be the number of times it lands inside a square, Y the number of heads. Assume $d = s/2$. Calculate:

c) $P(X = Y)$; d) $P(X < Y)$; e) $P(X > Y)$.

a) assume $d < s$

no difference if square coin side d .



$P(\text{coin in square})$
 $= P(0 < \text{right edge} < s-d,$
 $0 < \text{bot edge} < s-d)$
 $= \left(\frac{s-d}{s}\right)^2$

b) assume getting heads indep of position of coin. Assume fair coin.

$P(\text{coin in square, heads})$
 $= \frac{1}{2} \left(\frac{s-d}{s}\right)^2$

$$c) \quad d = s/2$$

$X = \# \text{ times (out of 4) coin lands in square}$

$$P(\text{coin in square}) = \left(\frac{s/2}{s}\right)^2 = \frac{1}{4}$$

$$X \sim \text{Bin}(4, 1/4)$$

$Y = \# \text{ heads (out of 4)}$

$$Y \sim \text{Bin}(4, 1/2)$$

$$P(X=Y) = P(X=0, Y=0) + P(X=1, Y=1)$$

$$+ P(X=2, Y=2) + P(X=3, Y=3) + P(X=4, Y=4)$$

$$= \binom{3}{4}^4 \cdot \left(\frac{1}{2}\right)^4 + 4^2 \frac{1}{4} \binom{3}{4}^3 \left(\frac{1}{2}\right)^4 + 6^2 \left(\frac{1}{4}\right)^2 \binom{3}{4}^2 \left(\frac{1}{2}\right)^4$$

$$+ 4^2 \left(\frac{1}{4}\right)^3 \binom{3}{4} \left(\frac{1}{2}\right)^4 + \left(\frac{1}{4}\right)^4 \left(\frac{1}{2}\right)^4$$

$$d) P(X < Y)$$

$$= P(X=0, Y=1) + P(X=0, Y=2) + P(X=0, Y=3) + P(X=0, Y=4)$$

$$+ P(X=1, Y=2) + P(X=1, Y=3) + P(X=1, Y=4)$$

$$+ P(X=2, Y=3) + P(X=2, Y=4) + P(X=3, Y=4)$$

This is straight-forward to find.

